

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 141 286 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
18.10.2006 Bulletin 2006/42

(51) Int Cl.:
C12N 15/12^(2006.01) **C07K 14/715^(2006.01)**
A61K 38/00^(2006.01) **A61K 39/00^(2006.01)**

(21) Application number: **99966166.3**

(86) International application number:
PCT/US1999/029493

(22) Date of filing: **13.12.1999**

(87) International publication number:
WO 2000/036103 (22.06.2000 Gazette 2000/25)

(54) **CYTOKINE RECEPTOR CHAIN**

CYTOKINREZEPTOR-KETTE

CHAINE DE RECEPTEURS DE CYTOKINE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(56) References cited:
EP-A- 0 812 913 **WO-A-97/31946**
WO-A-97/33913 **WO-A-97/47741**
WO-A-97/47742 **WO-A-98/10638**
WO-A1-97/15663 **FR-A- 2 742 156**
US-A- 5 599 905 **US-A- 5 696 234**

(30) Priority: **14.12.1998 US 211335**

(43) Date of publication of application:
10.10.2001 Bulletin 2001/41

(73) Proprietors:
• **Genetics Institute, LLC**
Cambridge, MA 02140 (US)
• **JOHNS HOPKINS UNIVERSITY**
Baltimore, MD 21218-2695 (US)

(72) Inventors:
• **COLLINS, Mary**
Natick, MA 01760 (US)
• **DONALDSON, Debra**
Medford, MA 02155 (US)
• **FITZ, Lori**
Arlington, MA 02174 (US)
• **NEBEN, Tamlyn**
Walnut Creek, CA 94598 (US)
• **WHITTERS, Matthew, J.**
Hudson, MA 01749 (US)
• **WOOD, Clive**
Boston, MA 02114 (US)
• **WILLS-KARP, Marsha,**
Johns Hopkins University
Baltimore, MD 21205 (US)

(74) Representative: **Dörries, Hans Ulrich et al**
Dörries, Frank-Molnia & Pohlman,
Triftstrasse 13
80538 München (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 141 286 B1

- CAPUT DANIEL ET AL: "Cloning and characterization of a specific interleukin (IL)-13 binding protein structurally related to the IL-5 receptor alpha chain." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 28, 1996, pages 16921-16926, XP002195292 ISSN: 0021-9258
- DONALDSON DEBRA D ET AL: "The murine IL-13 receptor alpha 2: molecular cloning, characterization, and comparison with murine IL-13 receptor alpha 1. " JOURNAL OF IMMUNOLOGY, vol. 161, no. 5, 1 August 1998 (1998-08-01), pages 2317-2324, XP002195293
- COLLINS M ET AL: "COMPARISON OF TWO DISTINCT LIGAND BINDING CHAINS FOR THE IL13 RECEPTOR IN THE MOUSE" JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, MOSBY - YEARLY BOOK, INC, US, vol. 99, no. 1, PART 2, 1997, page 287 XP000197588 ISSN: 0091-6749
- OBIRI ET AL.: 'Receptor for Interleukin 13: Interaction with interleukin 4 by a mechanism that does not involve the common gamma chain shared by receptors for interleukins 2, 4, 7, 9 and 15' THE JOURNAL OF BIOLOGICAL CHEMISTRY vol. 270, no. 15, 14 April 1995, pages 8797 - 8804, XP002927538
- ZURAWSKI ET AL.: 'The Primary Binding Subunit of the Human Interleukin-4 Receptor is Also a Component of the Interleukin-13 Receptor' THE JOURNAL OF BIOLOGICAL CHEMISTRY vol. 270, no. 23, 09 June 1995, pages 13869 - 13878, XP002927539
- VITA ET AL.: 'Characterization and Comparison of the Interleukin 13 Receptor with the Interleukin 4 Receptor on Several Cell Types' THE JOURNAL OF BIOLOGICAL CHEMISTRY vol. 270, no. 8, 24 February 1995, pages 3512 - 3517, XP002927540

DescriptionField of the Invention

[0001] The present invention relates to the use of an IL-13 antagonist in the manufacture of a medicament for treating an IL-13-related condition in a mammalian subject, wherein said antagonist is an antibody to IL-13 or an IL-13-binding fragment thereof and wherein said IL-13 related condition is atopy, allergic condition, asthma, immune complex disease, inflammatory condition of the lungs, immune deficiency, or cancer.

Background of the Invention

[0002] A variety of regulatory molecules, known as cytokines, have been identified including interleukin-13 (IL-13). Various protein forms of IL-13 and DNA encoding various forms of IL-13 activity are described in McKenzie et al., Proc. Natl. Acad. Sci. USA 90:3735 (1993); Minty et al., Nature 362:248 (1993); and Aversa et al., WO94/04680. Thus, the term "IL-13" includes proteins having the sequence and/or biological activity described in these documents, whether produced by recombinant genetic engineering techniques; purified from cell sources producing the factor naturally or upon induction with other factors; or synthesized by chemical techniques; or a combination of the foregoing.

[0003] IL-13 is a cytokine that has been implicated in production of several biological activities including: induction of IgG4 and IgE switching, including in human immature B cells (Punnonen et al., J. Immunol. 152:1094 (1994)); induction of germ line IgE heavy chain (ϵ) transcription and CD23 expression in normal human B cells (Punnonen et al., Proc. Natl. Acad. Sci. USA 90:3730 (1993)); and induction of B cell proliferation in the presence of CD40L or anti-CD40 mAb (Cocks et al., Int. Immunol. 5:657 (1993)). Although many activities of IL-13 are similar to those of IL-4, in contrast to IL-4, IL-13 does not have growth promoting effects on activated T cells or T cell clones (Zurawski et al., EMBO J. 12:2663 (1993)).

[0004] Like most cytokines, IL-13 exhibits certain biological activities by interacting with an IL-13 receptor ("IL-13R") on the surface of target cells. IL-13R and the IL-4 receptor ("IL-4R") sharing a common component, which is required for receptor activation; however, IL-13 does not bind to cells transfected with the 130 kD IL-4R (Zurawski et al., *supra*). Thus, the IL-13R must contain at least one other ligand binding chain. Cytokine receptors are commonly composed of two or three chains. The cloning of one ligand binding chain for IL-13 has been recently reported (Hilton et al., Proc. Natl. Acad. Sci. 93:497-501).

[0005] It would be desirable to identify and clone the sequence for any other IL-13 binding chain of IL-13R so that IL-13R proteins can be produced for various reasons, including production of therapeutics and screening for inhibitors of IL-13 binding to the receptor and receptor signaling.

Summary of the Invention

[0006] The present invention provides the use of an IL-13 antagonist in the manufacture of a medicament for treating an IL-13-related condition in a mammalian subject, wherein said antagonist is an antibody to IL-13 or an IL-13-binding fragment thereof and wherein said IL-13 related condition is atopy, allergic condition, asthma, immune complex disease, inflammatory condition of the lungs, immune deficiency, or cancer.

[0007] Another aspect of the invention is the use of an IL-13 antagonist in the manufacture of a medicament for inhibiting the interaction of IL-13 with an IL-13 binding protein in a condition selected from atopy, allergic condition, asthma, immune complex disease, inflammatory condition of the lungs, immune deficiency, or cancer, wherein said antagonist is an antibody to IL-13 or an IL-13-binding fragment thereof.

[0008] Also described are polynucleotides encoding the IL-13 binding chains of the interleukin-13 receptor are disclosed, including without limitation those from the murine and human receptors. In certain embodiments, the invention provides an isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO: 1 from nucleotide 256 to nucleotide 1404;
- (b) the nucleotide sequence of SEQ ID NO:3 from nucleotide 103 to nucleotide 1242;
- (c) a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (a) or (b) as a result of degeneracy of the genetic code;
- (d) a nucleotide sequence capable of hybridizing under stringent conditions to the nucleotide specified in (a) or (b);
- (e) a nucleotide sequence encoding a species homologue of the sequence specified in (a) or (b); and
- (f) an allelic variant of the nucleotide sequence specified in (a) or (b).

Preferably, the nucleotide sequence encodes a protein having a biological activity of the human IL-13 receptor. The nucleotide sequence may be operably linked to an expression control sequence. In preferred embodiments, the poly-

nucleotide comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 256 to nucleotide 1404; the nucleotide sequence of SEQ ID NO:1 from nucleotide 319 to nucleotide 1257; the nucleotide sequence of SEQ ID NO:1 from nucleotide 1324 to nucleotide 1404; the nucleotide sequence of SEQ ID NO:3 from nucleotide 103 to nucleotide 1242; the nucleotide sequence of SEQ ID NO:3 from nucleotide 178 to nucleotide 1125; or the nucleotide sequence of SEQ ID NO:3 from nucleotide 1189 to nucleotide 1242.

[0009] Further described are isolated polynucleotides comprising a nucleotide sequence encoding a peptide or protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding chain. Other preferred embodiments encode the amino acid sequence of SEQ ID NO:2 from amino acids 1 to 331 and the amino acid sequence of SEQ ID NO:2 from amino acids 26 to 331.

[0010] Host cells, preferably mammalian cells, transformed with the polynucleotides are also described.

[0011] Also described herein is a process for producing a IL-13bc protein. The process comprises:

- (a) growing a culture of the host cell of the present invention in a suitable culture medium; and
- (b) purifying the human IL-13bc protein from the culture.

Proteins produced according to these methods are also provided.

[0012] The Present invention also provides for an isolated IL-13bc protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having biological activity of the IL-13 receptor binding chain

Preferably the protein comprises the amino acid sequence of SEQ ID NO:2; the sequence from amino acid 22 to 334 of SEQ ID NO:2; the sequence of SEQ ID NO:4; or the sequence from amino acid 26 to 341 of SEQ ID NO:4. In other preferred embodiments, the specified amino acid sequence is part of a fusion protein (with an additional amino acid sequence not derived from IL-13bc). Preferred fusion proteins comprise an antibody fragment, such as an Fc fragment. Particularly preferred embodiments comprise the amino acid sequence of SEQ ID NO:2 from amino acids 1 to 331 and the amino acid sequence of SEQ ID NO:2 from amino acids 26 to 331.

[0013] Pharmaceutical compositions comprising a protein of the present invention and a pharmaceutically acceptable carrier are also described.

[0014] Further described are compositions comprising an antibody which specifically reacts with a protein as described herein.

[0015] Methods of identifying an inhibitor of IL-13 binding to the IL-13bc or IL-13 receptor are also described. These methods comprise:

- (a) combining an IL-13bc protein or a fragment thereof with IL-13 or a fragment thereof, said combination forming a first binding mixture;
- (b) measuring the amount of binding between the protein and the IL-13 or fragment in the first binding mixture;
- (c) combining a compound with the protein and the IL-13 or fragment to form a second binding mixture;
- (d) measuring the amount of binding in the second binding mixture; and
- (e) comparing the amount of binding in the first binding mixture with the amount of binding in the second binding mixture;

wherein the compound is capable of inhibiting IL-13 binding to the IL-13bc protein or IL-13 receptor when a decrease

in the amount of binding of the second binding mixture occurs. Inhibitors of IL-13R identified by these methods and pharmaceutical compositions containing them are also described.

[0016] Methods of inhibiting binding of IL-13 to the IL-13bc proteins or IL-13 receptor in a mammalian subject are also disclosed which comprise administering a therapeutically effective amount of a composition containing an IL-13bc protein,

[0017] Methods are also described for potentiating IL-13 activity, which comprise combining a protein having IL-13 activity with a protein of claim 11 and contacting such combination with a cell expressing at least one chain of IL-13R other than IL-13bc. Preferably, the contacting step is performed by administering a therapeutically effective amount of such combination to a mammalian subject.

[0018] Further methods are disclosed for treating an IL-13-tetated condition in a mammalian subject, said method comprising administering a therapeutically effective amount of a composition comprising an IL-13 antagonist and a pharmaceutically acceptable carrier. Other methods described relate to a method of inhibiting the interaction of IL-13 with an IL-13bc protein in a mammalian subject comprising administering a therapeutically effective amount of a composition comprising an IL-13 antagonist and a pharmaceutically acceptable carrier. Preferably, the antagonist is selected from the group consisting of an IL-13bc protein, a soluble form of IL-13R α 1, an antibody to IL-13 or an IL-13-binding fragment thereof, an antibody to IL-13bc or an IL-13bc-binding fragment thereof, an antibody to IL-13R α 1 or an IL-13R α 1-binding fragment thereof, IL-13R-binding mutants of IL-4, a small molecule capable of inhibiting the interaction of IL-13 with IL-13bc and a small molecule capable of inhibiting the interaction of IL-13 with IL-13R α 1.

Brief Description of the Figure

[0019]

Fig. 1: The figure presents photographs of IL-13, IL-4, IL-11 and mock transfected COS cells after exposure to IL-13bc-Fc as described in Example 4 below.

Fig. 2: Reversal of allergen-induced airway hyper responsiveness by in vivo blockade of interleukin-13. 10 days after initial intratracheal challenge, OVA- and PBS-immunized mice were again challenged intratracheally with either OVA or PBS. Mice were given sIL-13bc-Fc (400ug) or an equivalent amount of control hu-IgG by intraperitoneal injection on Day -1, 0, +1 and +3 of the secondary antigen challenge. The allergic phenotype was assessed 4 days after the PBS or OVA challenge. **(A)** Airway hyper responsiveness (AHR) to acetylcholine challenge, defined by the time-integrated rise in peak airway pressure (airway-pressure-time index [APT_I] in cmH₂O x sec). **(B)** Inflammatory cell composition of bronchoalveolar lavage fluids. Cell differential percentages were determined by light microscopic evaluation of cytopsin preparations. Data are expressed as absolute numbers of cells. **(C)** OVA-specific serum IgE concentrations. Results are means \pm SEM of 8-10 animals per group. * P < 0.05 compared with respective PBS control groups; ** P < 0.05 compared to OVA/Hu-Ig group (one-way ANOVA followed by Fisher's least significant difference test for multiple comparisons).

Fig. 3: Effects of IL-13 blockade on allergen-driven increases in mucus-containing cells in the airway epithelium. Lung sections (N = 4 per experimental group, four sections per animal) were fixed in formalin, cut into 10um sections and stained with hematoxylin and eosin, and periodic acid Schiff. Representative sections are shown. Bars = 100 um. PBS/Hu-Ig: PBS-immunized and challenged controls, demonstrating few mucus-containing cells. OVA/Hu-Ig: allergen-induced increases in interstitial inflammatory cells and increases in the number of goblet cells containing mucus. OVA/sIL-13bc-Fc: dramatic inhibitory effect of IL-13 blockade on allergen-induced goblet cell mucus production.

Fig. 4: IL-13 induction of airway hyperreactivity. Naive mice were given recombinant IL-13 (5 ug/mouse, 50 ul volume) or PBS daily by intratracheal instillation. 24 hrs after the last treatment, **(A)** Airway hyper responsiveness, **(B)** BAL eosinophil levels, **(C)** Serum total IgE levels, and **(D)** Mucus score were determined. Results are means \pm SEM (vertical bars) of 7-10 animals per group. * P < 0.05 compared to PBS group (Student's t test).

Detailed Description of Preferred Embodiments

[0020] The inventors of the present application have for the first time identified and provided polynucleotides encoding the IL-13 binding chain of IL-13R (hereinafter "IL-13bc"), including without limitation polynucleotides encoding murine and human IL-13bc.

[0021] SEQ ID NO: 1 provides the nucleotide sequence of a cDNA encoding the murine IL-13bc. SEQ ID NO:2 provides predicted the amino acid sequence of the receptor chain, including a putative signal sequence from amino acids 1-21. The mature murine IL-13bc is believed to have the sequence of amino acids 22-383 of SEQ ID NO:2. The mature murine receptor chain has at least three distinct domains: an extracellular domain (comprising approximately amino acids 22-334 of SEQ ID NO:2), a transmembrane domain (comprising approximately amino acids 335-356 of SEQ ID NO:2) and an

intracellular domain (comprising approximately amino acids 357-383 of SEQ ID NO:2).

[0022] SEQ ID NO:3 provides the nucleotide sequence of a cDNA encoding the human IL-13bc. SEQ ID NO:4 provides predicted the amino acid sequence of the receptor chain, including a putative signal sequence from amino acids 1-25. The mature human IL-13bc is believed to have the sequence of amino acids 26-380 of SEQ ID NO:4. The mature human receptor chain has at least three distinct domains: an extracellular domain (comprising approximately amino acids 26-341 of SEQ ID NO:4), a transmembrane domain (comprising approximately amino acids 342-362 of SEQ ID NO:4) and an intracellular domain (comprising approximately amino acids 363-380 of SEQ ID NO:4).

[0023] The first 81 amino acids of the human IL-13bc sequence are identical to the translated sequence of an expressed sequence tag (EST) identified as "yg99f10.r1 Homo sapiens cDNA clone 41648 5'" and assigned database accession number R52795.gb_est2. There are no homologies or sequence motifs in this EST sequence which would lead those skilled in the art to identify the encoded protein as a cytokine receptor. A cDNA clone corresponding to this database entry is publicly-available from the I.M.A.G.E. Consortium. Subsequent to the priority date of the present application, such clone was ordered by applicants and sequenced. The sequence of such clone was determined to be the sequence previously reported by applicants as SEQ ID NO:3 herein.

[0024] Soluble forms of IL-13bc protein can also be produced. Such soluble forms include without limitation proteins comprising amino acids 1-334 or 22-334 of SEQ ID NO:2 or amino acids 1-341 or 26-341 of SEQ ID NO:4. The soluble forms of the IL-13bc are further characterized by being soluble in aqueous solution, preferably at room temperature. IL-13bc proteins comprising only the intracellular domain or a portion thereof may also be produced. Any forms of IL-13bc of less than full length are referred to herein collectively with full length and mature forms as "IL-13bc" or "IL-13bc proteins." IL-13bc proteins of less than full length may be produced by expressing a corresponding fragment of the polynucleotide encoding the full-length IL-13bc protein (SEQ ID NO:1 or SEQ ID NO:3). These corresponding polynucleotide fragments are also described. Modified polynucleotides as described above may be made by standard molecular biology techniques, including construction of appropriate desired deletion mutants, site-directed mutagenesis methods or by the polymerase chain reaction using appropriate oligonucleotide primers.

[0025] A protein as described herein has "a biological activity of the IL-13 receptor binding chain" if it possess one or more of the following characteristics: (1) the ability to bind IL-13 or a fragment thereof (preferably a biologically active fragment thereof); and/or (2) the ability to interact with the second non-IL-13-binding chain of IL-13R to produce a signal characteristic of the binding of IL-13 to IL-13R. Preferably, the biological activity possessed by the protein is the ability to bind IL-13 or a fragment hereof, more preferably with a K_D of about 0.1 to about 100 nM. Methods for determining whether a particular protein or peptide has such activity include without limitation the methods described in the examples provided herein.

[0026] IL-13bc or active fragments thereof (IL-13bc proteins) may be fused to carrier molecules such as immunoglobulins. For example, soluble forms of the IL-13bc may be fused through "linker" sequences to the Fc portion of an immunoglobulin. Other fusions proteins, such as those with GST, Lex-A or MBP, may also be used.

[0027] Further described are allelic variants of the nucleotide sequences as set forth in SEQ ID NO:1 or SEQ ID NO:3, that is, naturally-occurring alternative forms of the isolated polynucleotide of SEQ ID NO:1 or SEQ ID NO:3 which also encode IL-13bc proteins, preferably those proteins having a biological activity of IL-13bc. Also described are isolated polynucleotides which hybridize to the nucleotide sequence set forth in SEQ ID NO:1 or SEQ ID NO:3 under highly stringent conditions (for example, 0.1xSSC at 65°C). Isolated polynucleotides which encode IL-13bc proteins but which differ from the nucleotide sequence set forth in SEQ ID NO:1 or SEQ ID NO:3 by virtue of the degeneracy of the genetic code are also described. Variations in the nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:3 which are caused by point mutations or by induced modifications are also described.

[0028] Also described are polynucleotides encoding homologues of the murine and human IL-13bc from other animal species, particularly other mammalian species. Species homologues can be identified and isolated by making probes or primers from the murine or human sequences disclosed herein and screening a library from an appropriate species, such as for example libraries constructed from PBMCs, thymus or testis of the relevant species.

[0029] The isolated polynucleotides may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al., *Nucleic Acids Res.* **19**, 4485-4490 (1991), in order to produce the IL-13bc protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* **185**, 537-566 (1990). As defined herein "operably linked" means enzymatically or chemically ligated to form a covalent bond between the isolated polynucleotide of the invention and the expression control sequence, in such a way that the IL-13bc protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

[0030] A number of types of cells may act as suitable host cells for expression of the IL-13bc protein. Any cell type capable of expressing functional IL-13bc protein may be used. Suitable mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from

in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK. HL-60, U937, HaK, Rat2, BaF3, 32D, FDCP-1, PC12, Mix or C2C12 cells.

[0031] The IL-13bc protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987). Soluble forms of the IL-13bc protein may also be produced in insect cells using appropriate isolated polynucleotides as described above.

[0032] Alternatively, the IL-13bc protein may be produced in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins.

[0033] Expression in bacteria may result in formation of inclusion bodies incorporating the recombinant protein. Thus, refolding of the recombinant protein may be required in order to produce active or more active material. Several methods for obtaining correctly folded heterologous proteins from bacterial inclusion bodies are known in the art. These methods generally involve solubilizing the protein from the inclusion bodies, then denaturing the protein completely using a chaotropic agent. When cysteine residues are present in the primary amino acid sequence of the protein, it is often necessary to accomplish the refolding in an environment which allows correct formation of disulfide bonds (a redox system). General methods of refolding are disclosed in Kohno, Meth. Enzym., 185:187-195 (1990). EP 0433225 and copending application USSN 08/163,877 describe other appropriate methods.

[0034] The IL-13bc protein may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a polynucleotide sequence encoding the IL-13bc protein.

[0035] The IL-13bc protein may be prepared by growing a culture transformed host cells under culture conditions necessary to express the desired protein. The resulting expressed protein may then be purified from the culture medium or cell extracts. Soluble forms of the IL-13bc protein can be purified from conditioned media. Membrane-bound forms of IL-13bc protein can be purified by preparing a total membrane fraction from the expressing cell and extracting the membranes with a non-ionic detergent such as Triton X-100.

[0036] The IL-13bc protein can be purified using methods known to those skilled in the art. For example, the IL-13bc protein can be concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) or polyethyleimine (PEI) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred (e.g., S-Sepharose® columns). The purification of the IL-13bc protein from culture supernatant may also include one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; or by hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or by immunoaffinity chromatography. Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the IL-13bc protein. Affinity columns including IL-13 or fragments thereof or including antibodies to the IL-13bc protein can also be used in purification in accordance with known methods. Some or all of the foregoing purification steps, in various combinations or with other known methods, can also be employed to provide a substantially purified isolated recombinant protein. Preferably, the isolated IL-13bc protein is purified so that it is substantially free of other mammalian proteins.

[0037] IL-13bc proteins may also be used to screen for agents which are capable of binding to IL-13bc or IL-13R or which interfere with the binding of IL-13 to the IL-13 or IL-13bc (either the extracellular or intracellular domains) and thus may act as inhibitors of normal binding and cytokine action ("IL-13R inhibitors"). Binding assays using a desired binding protein, immobilized or not, are well known in the art and may be used for this purpose using the IL-13bc protein of the invention. Purified cell based or protein based (cell free) screening assays may be used to identify such agents. For example, IL-13bc protein may be immobilized in purified form on a carrier and binding to purified IL-13bc protein may be measured in the presence and in the absence of potential inhibiting agents. A suitable binding assay may alternatively employ a soluble form of IL-13bc of the invention. Another example of a system in which inhibitors may be screened is described in Example 2 below.

[0038] In such a screening assay, a first binding mixture is formed by combining IL-13 or a fragment thereof and IL-13bc protein, and the amount of binding in the first binding mixture (B_0) is measured. A second binding mixture is also formed by combining IL-13 or a fragment thereof, IL-13bc protein, and the compound or agent to be screened, and the

amount of binding in the second binding mixture (B) is measured. The amounts of binding in the first and second binding mixtures are compared, for example, by performing a calculation of the ratio B/B_0 . A compound or agent is considered to be capable of inhibiting binding if a decrease in binding in the second binding mixture as compared to the first binding mixture is observed. Optionally, the second chain of IL-13R can be added to one or both of the binding mixtures. The formulation and optimization of binding mixtures is within the level of skill in the art, such binding mixtures may also contain buffers and salts necessary to enhance or to optimize binding, and additional control assays may be included in the screening assay of the invention.

[0039] Compounds found to reduce the binding activity of IL-13bc protein to IL-13 or its fragment to any degree, preferably by at least about 10%, more preferably greater than about 50% or more, may thus be identified and then secondarily screened in other binding assays and in vivo assays. By these means compounds having inhibitory activity for IL-13bc binding which may be suitable as therapeutic agents may be identified.

[0040] IL-13bc proteins, and polynucleotides encoding them, may also be used as diagnostic agents for detecting the expression or presence of IL-13bc, IL-13R, IL-13 or cells expressing IL-13bc, IL-13R or IL-13. The proteins or polynucleotides may be employed for such purpose in standard procedures for diagnostics assays using these types of materials. Suitable methods are well known to those skilled in the art.

[0041] As used herein "IL-13R" refers to IL-13bc and/or a second IL-13 receptor chain known as "IL-13R α 1" or "NR4" (see: murine receptor chain, Hilton et al., Proc. Natl. Acad. Sci. USA 1996,93:497-501; human receptor chain, Aman et al., J. Biol. Chem. 1996, 271:29265-70, and Gauchat et al., Eur. J. Immunol. 1997, 27:971-8).

[0042] IL-13bc acts as a mediator of the known biological activities of IL-13. As a result, IL-13bc protein (particularly, soluble IL-13bc proteins), IL-13R inhibitors (i.e., antagonists of interaction of IL-13 with IL-13R (such as, for example, antibodies to IL-13R (including particularly to IL-13bc or to IL-13R α 1) and fragments thereof, antibodies to IL-13 and fragments thereof, soluble IL-13R α 1 proteins, and small molecule and other inhibitors of the interaction of IL-13 with IL-13R (including with IL-13bc and/or with IL-13R α 1) may be useful in treatment or modulation of various medical conditions in which IL-13 is implicated or which are effected by the activity (or lack thereof) of IL-13 (collectively "IL-13-related conditions"). Mutated forms of IL-4 which bind to IL-13R can also be used as IL-13 antagonists (see, for example, those disclosed in Shanafelt et al., Proc. Natl. Acad. Sci. USA 1998, 95:9454-8; Aversa et al., J. Exp. Med. 1993, 178:2213-8; and Grunewald et al., J. Immunol. 1998, 160:4004-9).

[0043] IL-13-related conditions include without limitation Ig-mediated conditions and diseases, particularly IgE-mediated conditions (including without limitation atopy, allergic conditions, asthma, immune complex diseases (such as, for example, lupus, nephrotic syndrome, nephritis, glomerulonephritis, thyroiditis and Grave's disease)); inflammatory conditions of the lungs; immune deficiencies, specifically deficiencies in hematopoietic progenitor cells, or disorders relating thereto; cancer and other disease. Such pathological states may result from disease, exposure to radiation or drugs, and include, for example, leukopenia, bacterial and viral infections, anemia, B cell or T cell deficiencies such as immune cell or hematopoietic cell deficiency following a bone marrow transplantation. Since IL-13 inhibits macrophage activation, IL-13bc proteins may also be useful to enhance macrophage activation (i.e., in vaccination, treatment of mycobacterial or intracellular organisms, or parasitic infections).

[0044] IL-13bc proteins may also be used to potentiate the effects of IL-13 in vitro and in vivo. For example, an IL-13bc protein can be combined with a protein having IL-13 activity (preferably IL-13) and the resulting combination can be contacted with a cell expressing at least one chain of IL-13R other than IL-13bc (preferably all chains of IL-13R other than IL-13bc, such as IL-13R α 1). Preferably, the contacting step is performed by administering a therapeutically effective amount of such combination to a mammalian subject in vivo. The pre-established association of the IL-13 protein with the IL-13bc protein will aid in formation of the complete IL-13/IL-13R complex necessary for proper signaling. See for example the methods described by Economides et al., Science 270:1351 (1995).

[0045] IL-13bc protein and IL-13R inhibitors, purified from cells or recombinantly produced, may be used as a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may contain, in addition to IL-13bc or inhibitor and carrier, various diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration.

[0046] The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-14, IL-15, G-CSF, stem cell factor, and erythropoietin. The pharmaceutical composition may also include anti-cytokine antibodies. The pharmaceutical composition may contain thrombolytic or anti-thrombotic factors such as plasminogen activator and Factor VIII. The pharmaceutical composition may further contain other anti-inflammatory agents. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with isolated IL-13bc protein or IL-13bc inhibitor, or to minimize side effects caused by the isolated IL-13bc or IL-13bc inhibitor. Conversely, isolated IL-13bc or IL-13bc inhibitor may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of

the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

[0047] The pharmaceutical composition of the invention may be in the form of a liposome in which isolated IL-13bc protein or IL-13bc inhibitor is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers which in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

[0048] As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, e.g., amelioration of symptoms of, healing of, or increase in rate of healing of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

[0049] In practicing the method of treatment or use of the present invention, a therapeutically effective amount of isolated IL-13bc protein or IL-13bc inhibitor is administered to a mammal. Isolated IL-13bc protein or IL-13bc inhibitor may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, IL-13bc protein or IL-13bc inhibitor may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering IL-13bc protein or IL-13bc inhibitor in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

[0050] Administration of IL-13bc protein or IL-13bc inhibitor used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. Intravenous administration to the patient is preferred.

[0051] When a therapeutically effective amount of IL-13bc protein or IL-13bc inhibitor is administered orally, IL-13bc protein or IL-13bc inhibitor will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% IL-13bc protein or IL-13bc inhibitor, and preferably from about 25 to 90% IL-13bc protein or IL-13bc inhibitor. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of IL-13bc protein or IL-13bc inhibitor, and preferably from about 1 to 50% IL-13bc protein or IL-13bc inhibitor.

[0052] When a therapeutically effective amount of IL-13bc protein or IL-13bc inhibitor is administered by intravenous, cutaneous or subcutaneous injection, IL-13bc protein or IL-13bc inhibitor will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to IL-13bc protein or IL-13bc inhibitor an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additive known to those of skill in the art.

[0053] The amount of IL-13bc protein or IL-13bc inhibitor in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of IL-13bc protein or IL-13bc inhibitor with which to treat each individual patient. Initially, the attending physician will administer low doses of IL-13bc protein or IL-13bc inhibitor and observe the patient's response. Larger doses of IL-13bc protein or IL-13bc inhibitor may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not generally increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.1 μ g to about 100 mg of IL-13bc protein or IL-13bc inhibitor per kg body weight.

[0054] The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the IL-13bc protein or IL-13bc inhibitor will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

[0055] IL-13bc proteins of the invention may also be used to immunize animals to obtain polyclonal and monoclonal

antibodies which specifically react with the IL-13bc protein and which may inhibit binding of IL-13 or fragments thereof to the receptor. Such antibodies may be obtained using the entire IL-13bc as an immunogen, or by using fragments of IL-13bc, such as the soluble mature IL-13bc. Smaller fragments of the IL-13bc may also be used to immunize animals. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Additional peptide immunogens may be generated by replacing tyrosine residues with sulfated tyrosine residues. Methods for synthesizing such peptides are known in the art, for example, as in R.P. Merrifield, J.Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, et al., FEBS Lett. 211, 10 (1987).

[0056] Neutralizing or non-neutralizing antibodies (preferably monoclonal antibodies) binding to IL-13bc protein may also be useful therapeutics for certain tumors and also in the treatment of conditions described above. These neutralizing monoclonal antibodies may be capable of blocking IL-13 binding to the IL-13bc.

Example 1

Isolation of IL-13bc cDNAs

Isolation of the murine IL-13 receptor chain.

[0057] 5 ug of polyA⁺ RNA was prepared from the thymuses of 6-8 week old C3H/HeJ mice. Double stranded, hemimethylated cDNA was prepared using Stratagene's cDNA synthesis kit according to manufacturers instructions. Briefly, the first strand was primed with an oligodT-Xho primer, and after second strand synthesis, EcoRI adapters were added, and the cDNA was digested with XhoI, and purified. The cDNA was ligated to the XhoI-EcoRI sites of the Zap Express (Stratagene) lambda vector, and packaged using Gigapak II Gold packaging extracts (Stratagene) according to the manufacturers instructions. A library of 1.5×10^6 resulting recombinant phage was amplified following manufacturer's instructions. This library was screened with a degenerate 17mer oligonucleotide probe of the sequence KSRCTCCABK CRCTCCA (SEQ ID NO:5) (K = G+T; S= C+G; R=A+G; B=C+G+T) using standard TMAC hybridization conditions as described (Current Protocols in Molecular Biology, Ausubel, et al., editors., John Wiley and Sons, 1995, section 6.4.3). Clone A25 was identified because it hybridized to the 17mer probe, but not to probes derived from known hematopoietin receptors. This clone was isolated in plasmid form from the ZapExpress vector as per manufacturers instruction, and the DNA sequence was determined. The DNA sequence encoded a novel member of the hematopoietin receptor family.

[0058] Clone A25 containing the polynucleotide having the sequence of SEQ ID NO:1 was deposited with ATCC as pA25pBKCMV at accession number 69997 on February 22, 1996.

Isolation of the human IL-13 receptor chain.

[0059] A partial fragment of the human homolog of the murine receptor was isolated by PCR using oligonucleotides derived from the murine sequence. cDNA was prepared from human testis polyA⁺ RNA that was obtained from Clontech. A DNA fragment of 274 base pairs was amplified from this cDNA by PCR with the following oligonucleotides: ATAGT-TAAACCATGCGCACC (SEQ ID NO:6) and CTCCATTCGCTCCAAATTCC (SEQ ID NO:7) using AmpliTaq polymerase (Promega) in 1X Taq buffer containing 1.5 mM MgCl₂ for 30 cycles of incubation (94 °C x 1 minute, 42 °C for 1 minute, and 72 °C for 1 minute). The DNA sequence of this fragment was determined, and two oligonucleotides were prepared from an internal portion of this fragment with the following sequence: AGTCTATCTTACTITTACTCG (SEQ ID NO:8) and CATCTGAGCAATAAATATTCAC (SEQ ID NO:9). These oligonucleotides were used as probes to screen a human testis cDNA library purchased from CLONTECH (cat #HL1161). Filters were hybridized at 52 °C using standard 5XSSC hybridization conditions and washed in 2X SSC at 52 °C. Twenty two clones were isolated that hybridized to both oligonucleotides in a screen of 400,000 clones. DNA sequence was determined from four of the cDNA clones, and all encoded the same novel hematopoietin receptor. The predicted DNA sequence of the full length human receptor chain is shown as SEQ ID NO:3.

[0060] The human clone was deposited with ATCC as pH25#11pDR2 at accession number 69998 on February 22, 1996.

Example 2

Expression of Soluble IL-13bc Protein and Assay of Activity

Production and purification of soluble IL-13bc-Ig.

[0061] DNA encoding amino acids 1-331 of the extracellular domain of murine IL-13bc was fused to a spacer sequence

EP 1 141 286 B1

encoding gly-ser-gly by PCR and ligated in frame with sequences encoding the hinge CH2 CH3 regions of human IgG1 of the COS-1 expression vector pED.Fc. IL-13bc-Ig was produced from DEAE-dextran transfected COS-1 cells and purified via protein A sepharose chromatography (Pharmacia).

B9 proliferation assay

[0062] Stimulation of proliferation of B9 cells (Aarden et al. Eur. J. Immunol. 1987. 17:1411-1416) in response to IL-13 or IL-4 was measured by 3H-thymidine incorporation into DNA. Cells (5 x 10³/well) were seeded into 96 well plates with media containing growth factors at varying concentrations in the presence or absence of IL-13bc-Ig at 1ug/ml. After incubation for 3 days 1uCi/well of 3H-thymidine was added and the cells incubated for an additional 4 hrs. Incorporated radioactivity was determined using a LKB 1205 Plate reader.

[0063] The B9 cell line proliferated in response to IL-13, IL-4 or IL-6. Only responses to IL-13 were inhibited by the soluble IL-13bc-Ig, indicating that this receptor binds IL-13 specifically, but not IL-4 or IL-6. The tables show cpm. Two separate experiments are shown.

cytokine dilution	IL-13 (3ng/ml)	IL-13 plus A25-Fc (1ug/ml)	IL-4 (20 ng/ml)	IL-4 plus A25-Fc (1 ug/ml)	Cos IL-6 (1/10,000)
1	37734	1943	6443	6945	37887
1/3	30398	1571	2680	2442	36500
1/10	16101	1461	1767	1771	33335
1/30	2148	1567	1619	1783	27271
1/100	1574	1419	1522	1576	18831
1/300	1512	1531	1373	1577	7768
1/1000	1316	1392	1190	1474	2760
1/3000	1834	1994	1482	1819	1672

cytokine dilution	IL-13 (3ng/ml)	IL-13 plus A25-Fc (5ug/ml)	IL-4 (20 ng/ml)	IL-4 plus A25-Fc (5ug/ml)	Cos IL-6 (1/10,000)	Cos IL-6 plus A25-Fc (5ug/ml)
1	6413	295	1216	1158	6969	7703
1/3	5432	281	518	656	7827	8804
1/10	2051	281	489	520	8345	10027
1/30	506	319	279	476	8680	9114
1/100	430	372	288	423	7426	10364
1/300	330	287	323	420	5531	6254
1/1000	326	389	348	nt	2524	nt
no cytokine	339	279	404	394	326	279

Example 3

Direct Binding of Soluble IL-13bc to IL-13 Measured by Surface Plasmon Resonance (Biacore Analysis).

[0064] A Biacore biosensor was used to measure directly the specific binding of IL-13 to purified IL-13bc-Ig (Pharmacia, Johnsson et al., 1991). Approximately 10,000 to 17,000 resonance units (RU) of purified IL-13bc-Ig, human IgG1 or irrelevant receptor were each covalently immobilized to different flow cells on the sensor chip as recommended by the manufacturer. (RU's are a reflection of the mass of protein bound to the sensor chip surface.) Purified IL-13 was injected across the flow cells at 5 ul/min for 10 mins in the presence or absence of excess purified IL-13bc-Ig. Binding was quantified as the difference in RU before and after sample injection. Specific IL-13 binding of 481.9 RU was observed

EP 1 141 286 B1

only for immobilized IL-13bc-Ig whereas coinjection of IL-13 plus IL-13bc-Ig resulted in no binding to the immobilized IL-13bc-Ig (4 RU). No IL-13 binding was observed for either immobilized IgG or IL-11R-Ig (5.4 and 3.7 RU respectively).

Sample	IL-13bc-Ig (10,383 RU)	IgG control (13,399 RU)	IL-11R-Ig (17,182 RU)
100 ng/ml human IL-13	481.9 RU bound	5.4 RU bound	3.7 RU bound
100 ng/ml human IL-13 + soluble IL-13bc-Ig	4.0 RU bound	not tested	not tested

Example 4

Binding of IL-13 Expressed in COS Cells to Labeled IL-13BC-Ig Fusion Protein: COS in situ Detection of IL-13 with IL-13bc-Fc

[0065] Expression vectors for IL-13, IL-4, IL-11 or empty vector were transfected into COS-1 cells in duplicated plates via the DEAE-dextran method. Two days after transfection cells were washed twice in phosphate buffered saline (PBS) and fixed in the culture dish for 10' at 4° C with methanol. Following fixation cells were washed twice with PBS then rinsed once with binding buffer (PBS, 1% (w/v) bovine serum albumin, 0.1% (w/v) sodium azide) and incubated for two hours at 4° C in binding buffer with IL-13bc-Fc at 1.0ug/ml or with relevant anti-cytokine antisera. Cells were washed twice with PBS and incubated at 40 C with shaking in alkaline phosphatase labeled Rabbit F(ab)2' anti-human IgG diluted 1:500 in binding buffer (for Fc fusion detection) or Rabbit F(ab)2' anti-rat IgG (for anti-cytokine detection). Cells were again washed twice in PBS. Alkaline phosphatase activity was visualized using nitro blue tetrazolium and 5-bromo-4-chloro-3-indolyl-phosphate.

[0066] Specific binding was visualized under the microscope. Only cells transfected with IL-13 showed specific binding to IL13bc-Ig. (see photo of transfected cells, the Figure).

Example 5

Other Systems for Determination Biological Activity of IL-13bc Protein

[0067] Other systems can be used to determine whether a specific IL-13bc protein exhibits a "biological activity" of IL-13bc as defined herein. The following are examples of such systems.

Assays for IL-13 Binding

[0068] The ability of a IL-13bc protein to bind IL-13 or a fragment thereof can be determine by any suitable assays which can detect such binding. Some suitable examples follow.

[0069] Binding of IL-13 to the extracellular region of the IL-13bc protein will specifically cause a rapid induction of phosphotyrosine on the receptor protein. Assays for ligand binding activity as measured by induction of phosphorylation are described below.

[0070] Alternatively, a IL-13bc protein (such as, for example, a soluble form of the extracellular domain) is produced and used to detect IL-13 binding. For example, a DNA construct is prepared in which the extracellular domain (truncated prior, preferably immediately prior, to the predicted transmembrane domain) is ligated in frame to a cDNA encoding the hinge C_H2 and C_H3 domains of a human immunoglobulin (Ig) γ 1. This construct is generated in an appropriate expression vector for COS cells, such as pED Δ C or pMT2. The plasmid is transiently transfected into COS cells. The secreted IL-13bc-Ig fusion protein is collected in the conditioned medium and purified by protein A chromatography.

[0071] The purified IL-13bc-Ig fusion protein is used to demonstrate IL-13 binding in a number of applications. IL-13 can be coated onto the surface of an enzyme-linked immunosorbent assay (ELISA) plate, and then additional binding sites blocked with bovine serum albumin or casein using standard ELISA buffers. The IL-13bc-Ig fusion protein is then bound to the solid-phase IL-13, and binding is detected with a secondary goat anti-human Ig conjugated to horseradish peroxidase. The activity of specifically bound enzyme can be measured with a colorimetric substrate, such as tetramethyl benzidine and absorbance readings.

[0072] IL-13 may also be expressed on the surface of cells, for example by providing a transmembrane domain or glucosyl phosphatidyl inositol (GPI) linkage. Cells expressing the membrane bound IL-13 can be identified using the IL-13bc-Ig fusion protein. The soluble IL-13bc-Ig fusion is bound to the surface of these cells and detected with goat anti-human Ig conjugated to a fluorochrome, such as fluorescein isothiocyanate and flow cytometry.

Interaction Trap

[0073] A yeast genetic selection method, the "interaction trap" [Gyuris et al, Cell 75:791-803, 1993], can be used to determine whether a IL-13bc protein has a biological activity of IL-13bc as defined herein. In this system, the expression of reporter genes from both LexAop-Leu2 and LexAop-LacZ relies on the interaction between the bait protein, for example in this case a species which interacts with human IL-13bc, and the prey, for example in this case the human IL-13bc protein. Thus, one can measure the strength of the interaction by the level of Leu2 or LacZ expression. The most simple method is to measure the activity of the LacZ encoded protein, β -galactosidase. This activity can be judged by the degree of blueness on the X-Gal containing medium or filter. For the quantitative measurement of β -galactosidase activity, standard assays can be found in "Methods in Yeast Genetics" Cold Spring Harbor, New York, 1990 (by Rose, M.D., Winston, F., and Hieter, P.).

[0074] In such methods, if one wishes to determine whether the IL-13bc protein interacts with a particular species (such as, for example, a cytosolic protein which binds to the intracellular domain of the IL-13bc *in vivo*), that species can be used as the "bait" in the interaction trap with the IL-13bc protein to be tested serving as the "prey", or *vice versa*.

Example 6

Treatment of Asthma Using Soluble IL-13bc Protein

[0075] A well-characterized murine model of allergic asthma was used, in which allergen exposure leads to airway hyper responsiveness ("AHR"), pulmonary eosinophilia, elevations in antigen-specific serum IgE levels, and increases in airway epithelial mucus content (3, 11). Male A/J mice were immunized intraperitoneally and subsequently challenged intratracheally with soluble ovalbumin (OVA), the allergic phenotype being assessed 4 days after antigen challenge (13). Blockade of IL-13 was performed by the systemic administration of a soluble IL-13bc-IgGfC fusion protein (sIL-13bc-Fc), which specifically binds to and neutralizes IL-13, 24 hours before subsequent intratracheal allergen challenge (14). Challenge of allergen-immunized mice resulted in significant increases in airway responsiveness to acetylcholine (15) (Fig. 2A). Blockade of IL-13 resulted in complete reversal of such established allergen-induced AHR; thus IL-13 is necessary for the expression of AHR in this model. The ability of IL-13 ablation to reverse AHR after full development of the phenotype of allergic asthma contrasts with the inability of IL-4 ablation to accomplish such a reversal. The mechanism underlying the effectiveness of IL-4R α blockade in reversing allergen-induced AHR may be the inhibition of IL-13-mediated processes, consistent with the fact that Stat6 activation is downstream of IL-4R α -mediated signaling for both cytokines. IL-13 is probably the primary CD4⁺ T cell-derived factor responsible for allergen-induced AHR.

[0076] To evaluate candidate mechanisms underlying IL-13-dependent expression of AHR, we characterized known allergic effector cascades. Eosinophils have been implicated as primary effector cells in asthma and asthmatic AHR (16), but inhibition of IL-13 prior to repeat antigen provocation did not significantly affect allergen-induced pulmonary eosinophilia (17) (Fig. 2B). To assess the relevance of IgE-mediated pathways, we measured OVA-specific serum IgE (18). OVA-specific levels of IgE were observed in OVA-sensitized and -challenged mice, whereas no antigen-specific antibody levels were detected in PBS-immunized and -challenged mice (Fig. 2C). Blockade of IL-13 did not alter OVA-specific IgE levels, a lack of suppression which is likely due to the fact that IL-13 blockade occurred after initial antigen priming and antibody formation. Nonetheless, these results show that AHR is not dependent upon IgE production in this model, consistent with reports that allergic AHR develops normally in IgE deficient and B cell deficient mice (19).

[0077] In congruence with the pathology of human asthma, allergic asthma in murine models is associated with a marked increase in the mucus content of the airway epithelium (5, 11). Mucus hypersecretion is particularly profound in autopsy specimens from patients who die of acute asthma attacks (20). Blockade of IL-13 reverses allergen-induced increases in mucus-containing cells in the airways (Fig. 3), demonstrating that allergen-induced increases in airway mucus content are dependent upon IL-13. IL-4 is also implicated in this process, as IL-4 transgenic mice display marked goblet cell hyperplasia in the absence of antigen sensitization (5). However, transfer of Th2 clones from both IL-4-deficient and control mice into murine airways induces mucus overproduction (21), suggesting, yet again, that the immunoregulatory role of IL-4 needs to be carefully differentiated from its role as an effector molecule.

[0078] Daily administration of recombinant IL-13 (rIL-13) to the airways of naive (unimmunized) mice induced AHR, demonstrating that increases in IL-13 activity were sufficient to induce AHR (Fig. 4A) (22). AHR developed by 72 hours after the start of rIL-13 administration. A significant influx of eosinophils into bronchoalveolar lavage fluid was observed early after rIL-13 administration, however pulmonary eosinophilia was not observed at the time of expression of AHR (Fig. 4B). Although the significance of the time course of eosinophil influx remains unclear, it suggests that IL-13 alone may be sufficient to initiate eosinophilic infiltration of the airways, perhaps through its ability to upregulate chemokine expression (23). Airway administration of rIL-13 also resulted in a time-dependent increase in total serum IgE (Fig. 4C) (24), in line with the previously-reported ability of IL-13 to regulate IgE synthesis (25). Increases in serum IgE were independent of any immunization with allergen, findings that resonate with the observation that the human asthmatic

phenotype correlates better with total, rather than allergen-specific, serum IgE concentrations (26). As predicted from the above IL-13 inhibition studies, the administration of rIL-13 induced an increase in airway mucus production (Fig. 4D) (27).

References and Notes

[0079]

1. R. M. Sly, *Ann. Allergy* **53**, 20 (1984); R. Evans et al., *Chest* **91**, 65S (1987); N. Halfon and P. W. Newcheck, *Am. J. Pub. Health* **76**, 1308 (1986); R. M. Jackson, M. R. Sears, R. Beaglehole, H. H. Rea, *Chest* **94**, 914 (1988); P. J. Gergen and K. B. Weiss, *JAMA* **264**, 1688 (1990); W. M. Vollmer, A. S. Buist, M. L. Osborne, *J. Clin. Epid.* **45**, 999 (1992).
2. R. Beasley, W. R. Roche, J. A. Roberts, S. T. Holgate, *Am. Rev. Respir. Dis.* **139**, 806 (1989); R. Pauwels, *Clin. Exp. Allergy* **19**, 395 (1989); J. Bousquet et al., *N. Eng. J. Med.* **323**, 1033 (1990).
3. S. H. Gavett et al., *Am. J. Resp. Cell. Mol. Biol.* **10**, 587 (1994); A. A. Gerblich, H. Salik, M. R. Schuyler, *Amer. Rev. Resp. Dis.* **143**, 533 (1991); C. J. Corrigan, A. B. Kay, *Am. Rev. Resp. Dis.* **141**, 970 (1990); D. S. Robinson et al., *N. Engl. J. Med.* **326**, 298 (1992); C. Walker et al., *Am. Rev. Resp. Dis.* **146**, 109 (1992); S. H. Gavett et al., *J. Exp. Med.* **182**, 1527 (1995); N. W. Lukacs, R. M. Strieter, S. W. Chensue, S. L. Kunkel, *Am. J. Resp. Cell Mol. Biol.* **10**, 526 (1994).
4. F. D. Finkelman et al., *J. Immunol.* **141**, 2335 (1988); J. M. Wang et al., *Eur. J. Immunol.* **19**, 701 (1989).
5. J. A. Rankin et al., *Proc. Natl Acad. Sci. USA* **93**, 7821 (1996).
6. G. Brusselle, J. Kips, G. Joos, H. Bluethmann, R. Pauwels, *Am. J. Resp. Cell. Mol. Biol.* **12**, 254 (1995); D. B. Corry, et al., *J. Exp. Med.* **183**, 109 (1996); P. S. Foster et al., *J. Exp. Med.* **183**, 195 (1996).
7. A. J. Coyle et al., *Am. J. Resp. Cell. Mol. Biol.* **13**, 54 (1995).
8. A. K. Abbas, K. M. Murphy, A. Sher, *Nature* **383**, 787 (1996).
9. S. P. Hogan, et al., *J. Immunol.* **161**, 1501 (1998).
10. J. Punnonen et al., *Proc. Natl. Acad. Sci. USA* **90**, 3730 (1993); R. de Waal Malefyt, C. G. Figdor, J. E. de Vries, *Res. Immunol.* **144**, 629 (1993); G. Zurawski, J. E. de Vries, *Immunol. Today* **15**, 19 (1994).
11. S. H. Gavett et al., *Am. J. Physiol.* **272**, L253 (1977); D. Kuperman, B. Schofield, M. Wills-Karp, M. J. Grusby, *J. Exp. Med.* **187**, 939 (1998).
12. S. M. Zurawski, G. Zurawski, *EMBO J.* **11**, 3905 (1993); S. M. Zurawski et al., *J. Biol. Chem.* **270**, (1995). J.-X. Lin et al., *Immunity* **2**, 331 (1995).
13. Six-week-old male A/J mice were obtained from The Jackson Laboratory (Bar Harbor, ME) and were housed under laminar flow hoods in an environmentally-controlled specific pathogen-free animal facility for the duration of experiments (N = 4-10 mice/experimental group). The studies reported here conformed to the principles for laboratory animal research outlined by the Animal Welfare Act and the Department of Health, Education and Welfare (N.I.H.) guidelines for the experimental use of animals. Mice were immunized by an intraperitoneal injection of 10 ug ovalbumin (OVA; Crude grade IV, Sigma; St. Louis, MO) in 0.2 ml PBS or PBS alone. 14 days after immunization, mice were anesthetized with a mixture of ketamine and xylazine (45 and 8 mg/kg, respectively) and challenged intratracheally with 50 ul of a 1.5% solution of OVA or an equivalent volume of PBS as a control. 10 days after this first antigen challenge, mice were challenged again intratracheally with either OVA or PBS. Characterization of the allergic phenotype was performed 96 hours after the second antigen challenge.
14. Human IL-13bc was cloned as described above. For soluble expression of the murine homolog, a pED expression vector containing DNA encoding the murine sIL-13bc extracellular domain, fused in frame with the hinge CH2/CH3 regions of human IgG1 (as described in previous examples), was transfected into CHO cells [D. D. Donaldson et al., *J. Immunol.* **161**, 2317 (1998)]. The sIL-13bc-Fc was purified with rProtein A-Sepharose [J. F. Urban et al., *Immunity* **8**, 255 (1998)]. The in vitro ID₅₀, as determined by the ability to neutralize 3 ng/ml of murine IL-13 in the B9 proliferation assay was approximately 10 ng/ml. Human IgG, used as a control for sIL-13bc-Fc, was similarly purified by rProtein A-Sepharose chromatography from a 10% solution of human immune globulin that is commercially available for intravenous administration (Miles) [ibid]. Mice were given sIL-13bc-Fc (400ug), or an equivalent amount of the control hu-IgG, by intraperitoneal injection on Day -1, 0, +1, and +3 of secondary antigen challenge.
15. Airway reactivity to intravenous administration of acetylcholine was measured (11), 3 days after final intratracheal challenge. Mice were anesthetized with sodium pentobarbital (90 mg/kg), intubated, ventilated at a rate of 120 breaths/minute with a constant tidal volume of air (0.2 ml), and paralyzed with decamethonium bromide (25 mg/kg). After establishment of a stable airway pressure, acetylcholine was injected intravenously (50 ug/kg) and dynamic airway pressure was followed for 5 minutes.
16. G. J. Gleich, *J. All. Clin. Immunol.* **8**, 422 (1990).
17. Bronchoalveolar lavage was conducted as described (11).

18. A kidney was excised, and pooled blood was collected for antibody analysis as described (11). Serum was separated by centrifugation and stored at -80°C until analysis. Serum OVA-specific IgE levels were determined by sandwich ELISA. Sample wells were coated with a 0.01% OVA solution in PBS, blocked with 10% FBS in PBS, and washed with 0.05% Tween-20 in PBS. Serum samples were diluted 1:10 and 1:100 with 10% FBS in PBS. After an overnight incubation, plates were washed with 0.05% Tween-20 in PBS and biotin-conjugated anti-mouse IgE (PharMingen, San Diego, CA) was added. After a wash, 0.0025 mg/ml avidin peroxidase (Sigma) in 10% FBS/PBS was added, and plates were developed with ABTS (2,2'-azino-bis[3-ethyl-benzthiazole sulfonate]) (Kirkegaard and Perry). Plates were read at 405 nm within 30 minutes. Reported O.D. values are of serum samples diluted 1:10 since these values were proven to be below the saturation point of the assay by comparison of O.D. values of serum samples diluted 1:100 with 10% FBS/PBS.

19. P. D. Mehlhop et al., *Proc. Natl. Acad. Sci. USA* **94**, 1344 (1997); M. Korsgren et al., *J. Exp. Med.* **185**, 885 (1997).

20. T. Aikawa et al., *Chest* **101**, 916 (1992).

21. L. Cohn, R. J. Homer, A. Marinov, J. Rankin, K. Bottomly, *J. Exp. Med.* **186**, 1737 (1997).

22. DNA encoding a honeybee melittin leader [D. C. Tessier, D. Y. Thomas, H. E. Khouri, F. Laliberte, T. Vernet, *Gene* **2**, 177 (1991)] followed by a six-histidine tag was fused by an enterokinase cleavage site to the mature region of murine IL-13 at Gly21 and constructed in the mammalian expression vector pHTop. H6-EK murine IL-13 protein was produced from stably-transfected CHO cells and purified via Ni-NTA chromatography to greater than 97% purity as determined by SDS-PAGE. Protein concentration was determined by absorption at 280 nm and endotoxin contamination was less than 30 EU/mg as measured by Cape Cod Associates LAL assay. The ED₅₀ of H6-EK murine IL-13 as determined by the Ba/F3.IL-13R 1 proliferation assay was 1 ng/ml. Murine rIL-13 (5 µg in a total volume of 50 µl) was administered daily by intratracheal instillation to naive mice anesthetized with a mixture of ketamine and xylazine (45 and 8 mg/kg, respectively).

23. M. Goebeler et al., *Immunol.* **91**, 450 (1997).

24. A murine IgE-specific ELISA was used to quantitate total IgE immunoglobulin levels in serum using complementary antibody pairs for mouse IgE (R35-72 and R35-92) obtained from PharMingen according to the manufacturer's instructions. Duplicate samples (of a 1/10 dilution in 10% FBS in PBS) were examined from each animal. O.D. readings of samples were converted to pg/ml using values obtained from standard curves generated with known concentrations of recombinant mouse IgE (5-2000 pg/ml), and the final concentration was obtained by multiplying by the dilution factor.

25. C. L. Emson, S. E. Bell, A. Jones, W. Wisden, A. N. J. McKeazie, *J. Exp. Med.* **188**, 399 (1998).

26. L. R. Friedhoff, D. G. Marsh, *Int. Arch. All. Immunol.* **100**, 355 (1993).

27. To examine the effects of rIL-13 on mucus cell content of the airway epithelium, lungs were excised and fixed in 10% formalin. They were then washed in 70% ethanol, dehydrated, embedded in glycol methacrylate, cut into 10 µm sections, mounted on slides, and stained with hematoxylin and eosin and periodic acid Schiff. Four sections were examined per animal; 4 fields were scored per lung section. Sections were scored on a scale from 1-4 with 1 representing no mucus cell content.

28. J. Luyimbazi, X. Xu, M. Wills-Karp, unpublished results.

29. C. Walker et al., *Am. Rev. Respir. Dis.* **146**, 109 (1992); M. Humbert et al., *J. All. Clin. Immunol.* **99**, 657 (1997); S. K. Huang, *J. Immunol.* **155**, 2688 (1995).

30. D. G. Marsh, et al., *Science* **264**, 1152 (1996).

31. L. J. Rosenwasser, *N. Engl. J. Med.* **337**, 1766 (1977).

32. G. K. Hershey et al., *N Engl. J. Med.* **337**, 1720 (1997).

SEQUENCE LISTING

[0080]

<110> Collins, Mary
Donaldson, Debra
Fitz, Lora
Neben, Tamlyn
Whitters, Matthew
Wood, Clive
Wills-Karp, Marsha
Genetics Institute, Inc.
Johns Hopkins University
<120> Cytokine Receptor Chain

EP 1 141 286 B1

<130> GI 5268A

<140>

<141>

<160> 9

5 <170> PatentIn Ver. 2.0

<210> 1

<211> 1525

<212> DNA

<213> Mus sp.

10 <220>

<221> CDS

<222> (256)..(1404)

<400> 1

```

15      gaattcggca cgaggggagag gaggagggaa agatagaaag agagagagaa agattgcttg 60
      ctaccctga acagtgacct ctctcaagac agtgctttgc tcttcacgta taaggaagga 120
      aaacagtaga gattcaattt agtgtctaata gtggaaagga ggacaaagag gtcttgatgat 180
20      aactgcctgt gataatacat ttcttgagaa accatattat tgagtagagc tttcagcaca 240
      ctaaactctg gagaa atg gct ttt gtg cat atc aga tgc ttg tgt ttc att 291
      Met Ala Phe Val His Ile Arg Cys Leu Cys Phe Ile
      1 5 10
25      ctt ctt tgt aca ata act ggc tat tct ttg gag ata aaa gtt aat cct 339
      Leu Leu Cys Thr Ile Thr Gly Tyr Ser Leu Glu Ile Lys Val Asn Pro
      15 20 25
30      cct cag gat ttt gaa ata ttg gat cct gga tta ctt ggt tat ctc tat 387
      Pro Gln Asp Phe Glu Ile Leu Asp Pro Gly Leu Leu Gly Tyr Leu Tyr
      30 35 40
35      ttg caa tgg aaa cct cct gtg gtt ata gaa aaa ttt aag ggc tgt aca 435
      Leu Gln Trp Lys Pro Pro Val Val Ile Glu Lys Phe Lys Gly Cys Thr
      45 50 55 60
40      cta gaa tat gag tta aaa tac cga aat gtt gat agc gac agc tgg aag 483
      Leu Glu Tyr Glu Leu Lys Tyr Arg Asn Val Asp Ser Asp Ser Trp Lys
      65 70 75

```

45

50

55

EP 1 141 286 B1

	act	ata	att	act	agg	aat	cta	att	tac	aag	gat	ggg	ttt	gat	ctt	aat	531
	Thr	Ile	Ile	Thr	Arg	Asn	Leu	Ile	Tyr	Lys	Asp	Gly	Phe	Asp	Leu	Asn	
				80					85					90			
5	aaa	ggc	att	gaa	gga	aag	ata	cgt	acg	cat	ttg	tca	gag	cat	tgt	aca	579
	Lys	Gly	Ile	Glu	Gly	Lys	Ile	Arg	Thr	His	Leu	Ser	Glu	His	Cys	Thr	
			95					100					105				
10	aat	gga	tca	gaa	gta	caa	agt	cca	tgg	ata	gaa	gct	tct	tat	ggg	ata	627
	Asn	Gly	Ser	Glu	Val	Gln	Ser	Pro	Trp	Ile	Glu	Ala	Ser	Tyr	Gly	Ile	
			110				115					120					
15	tca	gat	gaa	gga	agt	ttg	gaa	act	aaa	att	cag	gac	atg	aag	tgt	ata	675
	Ser	Asp	Glu	Gly	Ser	Leu	Glu	Thr	Lys	Ile	Gln	Asp	Met	Lys	Cys	Ile	
	125					130					135					140	
20	tat	tat	aac	tgg	cag	tat	ttg	gtc	tgc	tct	tgg	aaa	cct	ggc	aag	aca	723
	Tyr	Tyr	Asn	Trp	Gln	Tyr	Leu	Val	Cys	Ser	Trp	Lys	Pro	Gly	Lys	Thr	
				145					150						155		
25	gta	tat	tct	gat	acc	aac	tat	acc	atg	ttt	ttc	tgg	tat	gag	ggc	ttg	771
	Val	Tyr	Ser	Asp	Thr	Asn	Tyr	Thr	Met	Phe	Phe	Trp	Tyr	Glu	Gly	Leu	
				160					165					170			
30	gat	cat	gcc	tta	cag	tgt	gct	gat	tac	ctc	cag	cat	gat	gaa	aaa	aat	819
	Asp	His	Ala	Leu	Gln	Cys	Ala	Asp	Tyr	Leu	Gln	His	Asp	Glu	Lys	Asn	
			175					180					185				
35	gtt	gga	tgc	aaa	ctg	tcc	aac	ttg	gac	tca	tca	gac	tat	aaa	gat	ttt	867
	Val	Gly	Cys	Lys	Leu	Ser	Asn	Leu	Asp	Ser	Ser	Asp	Tyr	Lys	Asp	Phe	
		190					195					200					
40	ttt	atc	tgt	gtt	aat	gga	tct	tca	aag	ttg	gaa	ccc	atc	aga	tcc	agc	915
	Phe	Ile	Cys	Val	Asn	Gly	Ser	Ser	Lys	Leu	Glu	Pro	Ile	Arg	Ser	Ser	
	205					210					215					220	
45	tat	aca	gtt	ttt	caa	ctt	caa	aat	ata	gtt	aaa	cca	ttg	cca	cca	gaa	963
	Tyr	Thr	Val	Phe	Gln	Leu	Gln	Asn	Ile	Val	Lys	Pro	Leu	Pro	Pro	Glu	
				225					230						235		
50	ttc	ctt	cat	att	agt	gtg	gag	aat	tcc	att	gat	att	aga	atg	aaa	tgg	1011
	Phe	Leu	His	Ile	Ser	Val	Glu	Asn	Ser	Ile	Asp	Ile	Arg	Met	Lys	Trp	
				240					245					250			
55	agc	aca	cct	gga	gga	ccc	att	cca	cca	agg	tgt	tac	act	tat	gaa	att	1059
	Ser	Thr	Pro	Gly	Gly	Pro	Ile	Pro	Pro	Arg	Cys	Tyr	Thr	Tyr	Glu	Ile	
			255					260					265				
60	gtg	atc	cga	gaa	gac	gat	att	tcc	tgg	gag	tct	gcc	aca	gac	aaa	aac	1107
	Val	Ile	Arg	Glu	Asp	Asp	Ile	Ser	Trp	Glu	Ser	Ala	Thr	Asp	Lys	Asn	
		270					275					280					
65	gat	atg	aag	ttg	aag	agg	aga	gca	aat	gaa	agt	gaa	gac	cta	tgc	ttt	1155
	Asp	Met	Lys	Leu	Lys	Arg	Arg	Ala	Asn	Glu	Ser	Glu	Asp	Leu	Cys	Phe	
	285					290					295					300	
70	ttt	gta	aga	tgt	aag	gtc	aat	ata	tat	tgt	gca	gat	gat	gga	att	tgg	1203
	Phe	Val	Arg	Cys	Lys	Val	Asn	Ile	Tyr	Cys	Ala	Asp	Asp	Gly	Ile	Trp	
				305						310					315		

EP 1 141 286 B1

agc gaa tgg agt gaa gag gaa tgt tgg gaa ggt tac aca ggg cca gac 1251
 Ser Glu Trp Ser Glu Glu Glu Cys Trp Glu Gly Tyr Thr Gly Pro Asp
 320 325 330

5

tca aag att att ttc ata gta cca gtt tgt ctt ttc ttt ata ttc ctt 1299
 Ser Lys Ile Ile Phe Ile Val Pro Val Cys Leu Phe Phe Ile Phe Leu
 335 340 345

10

ttg tta ctt ctt tgc ctt att gtg gag aag gaa gaa cct gaa ccc aca 1347
 Leu Leu Leu Leu Cys Leu Ile Val Glu Lys Glu Glu Pro Glu Pro Thr
 350 355 360

15

ttg agc ctc cat gtg gat ctg aac aaa gaa gtg tgt gct tat gaa gat 1395
 Leu Ser Leu His Val Asp Leu Asn Lys Glu Val Cys Ala Tyr Glu Asp
 365 370 375 380

20

acc ctc tgt taaaccacca atttcttgac atagagccag ccagcaggag 1444
 Thr Leu Cys

tcatattaaa ctcaatttct cttaaaattt cgaatacatc ttcttgaaaa tccaaaaaaa 1504
 aaaaaaaaaa aaaaactcga g 1525

<210> 2
 <211> 383
 <212> PRT
 <213> Mus sp.
 <400> 2

30

Met Ala Phe Val His Ile Arg Cys Leu Cys Phe Ile Leu Leu Cys Thr
 1 5 10 15

35

Ile Thr Gly Tyr Ser Leu Glu Ile Lys Val Asn Pro Pro Gln Asp Phe
 20 25 30

Glu Ile Leu Asp Pro Gly Leu Leu Gly Tyr Leu Tyr Leu Gln Trp Lys
 35 40 45

40

Pro Pro Val Val Ile Glu Lys Phe Lys Gly Cys Thr Leu Glu Tyr Glu
 50 55 60

Leu Lys Tyr Arg Asn Val Asp Ser Asp Ser Trp Lys Thr Ile Ile Thr
 65 70 75 80

45

Arg Asn Leu Ile Tyr Lys Asp Gly Phe Asp Leu Asn Lys Gly Ile Glu
 85 90 95

Gly Lys Ile Arg Thr His Leu Ser Glu His Cys Thr Asn Gly Ser Glu
 100 105 110

50

Val Gln Ser Pro Trp Ile Glu Ala Ser Tyr Gly Ile Ser Asp Glu Gly
 115 120 125

Ser Leu Glu Thr Lys Ile Gln Asp Met Lys Cys Ile Tyr Tyr Asn Trp
 130 135 140

55

Gln Tyr Leu Val Cys Ser Trp Lys Pro Gly Lys Thr Val Tyr Ser Asp
 145 150 155 160

EP 1 141 286 B1

	Thr	Asn	Tyr	Thr	Met	Phe	Phe	Trp	Tyr	Glu	Gly	Leu	Asp	His	Ala	Leu
					165					170					175	
5	Gln	Cys	Ala	Asp	Tyr	Leu	Gln	His	Asp	Glu	Lys	Asn	Val	Gly	Cys	Lys
				180					185					190		
	Leu	Ser	Asn	Leu	Asp	Ser	Ser	Asp	Tyr	Lys	Asp	Phe	Phe	Ile	Cys	Val
			195					200					205			
10	Asn	Gly	Ser	Ser	Lys	Leu	Glu	Pro	Ile	Arg	Ser	Ser	Tyr	Thr	Val	Phe
		210					215					220				
	Gln	Leu	Gln	Asn	Ile	Val	Lys	Pro	Leu	Pro	Pro	Glu	Phe	Leu	His	Ile
15						230					235					240
	Ser	Val	Glu	Asn	Ser	Ile	Asp	Ile	Arg	Met	Lys	Trp	Ser	Thr	Pro	Gly
					245					250					255	
	Gly	Pro	Ile	Pro	Pro	Arg	Cys	Tyr	Thr	Tyr	Glu	Ile	Val	Ile	Arg	Glu
20				260					265					270		
	Asp	Asp	Ile	Ser	Trp	Glu	Ser	Ala	Thr	Asp	Lys	Asn	Asp	Met	Lys	Leu
			275					280					285			
25	Lys	Arg	Arg	Ala	Asn	Glu	Ser	Glu	Asp	Leu	Cys	Phe	Phe	Val	Arg	Cys
		290					295					300				
	Lys	Val	Asn	Ile	Tyr	Cys	Ala	Asp	Asp	Gly	Ile	Trp	Ser	Glu	Trp	Ser
		305				310					315					320
30	Glu	Glu	Glu	Cys	Trp	Glu	Gly	Tyr	Thr	Gly	Pro	Asp	Ser	Lys	Ile	Ile
					325					330					335	
	Phe	Ile	Val	Pro	Val	Cys	Leu	Phe	Phe	Ile	Phe	Leu	Leu	Leu	Leu	Leu
				340					345					350		
35	Cys	Leu	Ile	Val	Glu	Lys	Glu	Glu	Pro	Glu	Pro	Thr	Leu	Ser	Leu	His
			355					360					365			
	Val	Asp	Leu	Asn	Lys	Glu	Val	Cys	Ala	Tyr	Glu	Asp	Thr	Leu	Cys	
40		370					375					380				

<210> 3
 <211> 1369
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> CDS
 <222> (103)..(1245)
 <400> 3

EP 1 141 286 B1

ggatccgcgc ggatgaaggc tatttgaagt cgccataacc tggtcagaag tgtgcctgtc 60

ggcgggggaga gaggcaatat caaggtttta aatctcggag aa atg gct ttc gtt 114
Met Ala Phe Val

1

tgc ttg gct atc gga tgc tta tat acc ttt ctg ata agc aca aca ttt 162

EP 1 141 286 B1

	Cys	Leu	Ala	Ile	Gly	Cys	Leu	Tyr	Thr	Phe	Leu	Ile	Ser	Thr	Thr	Phe	
	5					10					15					20	
5	ggc	tgt	act	tca	tct	tca	gac	acc	gag	ata	aaa	ggt	aac	cct	cct	cag	210
	Gly	Cys	Thr	Ser	Ser	Ser	Asp	Thr	Glu	Ile	Lys	Val	Asn	Pro	Pro	Gln	
					25				30					35			
	gat	ttt	gag	ata	gtg	gat	ccc	gga	tac	tta	ggt	tat	ctc	tat	ttg	caa	258
10	Asp	Phe	Glu	Ile	Val	Asp	Pro	Gly	Tyr	Leu	Gly	Tyr	Leu	Tyr	Leu	Gln	
				40				45					50				
	tgg	caa	ccc	cca	ctg	tct	ctg	gat	cat	ttt	aag	gaa	tgc	aca	gtg	gaa	306
	Trp	Gln	Pro	Pro	Leu	Ser	Leu	Asp	His	Phe	Lys	Glu	Cys	Thr	Val	Glu	
			55				60					65					
15	tat	gaa	cta	aaa	tac	cga	aac	att	ggt	agt	gaa	aca	tgg	aag	acc	atc	354
	Tyr	Glu	Leu	Lys	Tyr	Arg	Asn	Ile	Gly	Ser	Glu	Thr	Trp	Lys	Thr	Ile	
		70				75					80						
	att	act	aag	aat	cta	cat	tac	aaa	gat	ggg	ttt	gat	ctt	aac	aag	ggc	402
20	Ile	Thr	Lys	Asn	Leu	His	Tyr	Lys	Asp	Gly	Phe	Asp	Leu	Asn	Lys	Gly	
		85				90				95						100	
	att	gaa	gcg	aag	ata	cac	acg	ctt	tta	cca	tgg	caa	tgc	aca	aat	gga	450
25	Ile	Glu	Ala	Lys	Ile	His	Thr	Leu	Leu	Pro	Trp	Gln	Cys	Thr	Asn	Gly	
				105				110							115		
	tca	gaa	ggt	caa	agt	tcc	tgg	gca	gaa	act	act	tat	tgg	ata	tca	cca	498
	Ser	Glu	Val	Gln	Ser	Ser	Trp	Ala	Glu	Thr	Thr	Tyr	Trp	Ile	Ser	Pro	
				120				125						130			
30	caa	gga	att	cca	gaa	act	aaa	ggt	cag	gat	atg	gat	tgc	gta	tat	tac	546
	Gln	Gly	Ile	Pro	Glu	Thr	Lys	Val	Gln	Asp	Met	Asp	Cys	Val	Tyr	Tyr	
			135				140						145				
	aat	tgg	caa	tat	tta	ctc	tgt	tct	tgg	aaa	cct	ggc	ata	ggt	gta	ctt	594
35	Asn	Trp	Gln	Tyr	Leu	Leu	Cys	Ser	Trp	Lys	Pro	Gly	Ile	Gly	Val	Leu	
		150				155					160						
	ctt	gat	acc	aat	tac	aac	ttg	ttt	tac	tgg	tat	gag	ggc	ttg	gat	cat	642
40	Leu	Asp	Thr	Asn	Tyr	Asn	Leu	Phe	Tyr	Trp	Tyr	Glu	Gly	Leu	Asp	His	
		165				170					175					180	
	gca	tta	cag	tgt	ggt	gat	tac	atc	aag	gct	gat	gga	caa	aat	ata	gga	690
	Ala	Leu	Gln	Cys	Val	Asp	Tyr	Ile	Lys	Ala	Asp	Gly	Gln	Asn	Ile	Gly	
				185				190							195		
45	tgc	aga	ttt	ccc	tat	ttg	gag	gca	tca	gac	tat	aaa	gat	ttc	tat	att	738
	Cys	Arg	Phe	Pro	Tyr	Leu	Glu	Ala	Ser	Asp	Tyr	Lys	Asp	Phe	Tyr	Ile	
				200				205						210			
	tgt	ggt	aat	gga	tca	tca	gag	aac	aag	cct	atc	aga	tcc	agt	tat	ttc	786
50	Cys	Val	Asn	Gly	Ser	Ser	Glu	Asn	Lys	Pro	Ile	Arg	Ser	Ser	Tyr	Phe	
			215					220					225				
	act	ttt	cag	ctt	caa	aat	ata	ggt	aaa	cct	ttg	cgg	cca	gtc	tat	ctt	834
	Thr	Phe	Gln	Leu	Gln	Asn	Ile	Val	Lys	Pro	Leu	Pro	Pro	Val	Tyr	Leu	
		230				235					240						
55	act	ttt	act	cgg	gag	agt	tca	tgt	gaa	att	aag	ctg	aaa	tgg	agc	ata	882

EP 1 141 286 B1

	Thr Phe Thr Arg Glu Ser Ser Cys Glu Ile Lys Leu Lys Trp Ser Ile	
	245 250 255 260	
5	cct ttg gga cct att cca gca agg tgt ttt gat tat gaa att gag atc 930	
	Pro Leu Gly Pro Ile Pro Ala Arg Cys Phe Asp Tyr Glu Ile Glu Ile	
	265 270 275	
10	aga gaa gat gat act acc ttg gtg act gct aca gtt gaa aat gaa aca 978	
	Arg Glu Asp Asp Thr Thr Leu Val Thr Ala Thr Val Glu Asn Glu Thr	
	280 285 290	
15	tac acc ttg aaa aca aca aat gaa acc cga caa tta tgc ttt gta gta 1026	
	Tyr Thr Leu Lys Thr Thr Asn Glu Thr Arg Gln Leu Cys Phe Val Val	
	295 300 305	
20	aga agc aaa gtg aat att tat tgc tca gat gac gga att tgg agt gag 1074	
	Arg Ser Lys Val Asn Ile Tyr Cys Ser Asp Asp Gly Ile Trp Ser Glu	
	310 315 320	
25	tgg agt gat aaa caa tgc tgg gaa ggt gaa gac cta tcg aag aaa act 1122	
	Trp Ser Asp Lys Gln Cys Trp Glu Gly Glu Asp Leu Ser Lys Lys Thr	
	325 330 335 340	
30	ttg cta cgt ttc tgg cta cca ttt ggt ttc atc tta ata tta gtt ata 1170	
	Leu Leu Arg Phe Trp Leu Pro Phe Gly Phe Ile Leu Ile Leu Val Ile	
	345 350 355	
35	ttt gta acc ggt ctg ctt ttg cgt aag cca aac acc tac cca aaa atg 1218	
	Phe Val Thr Gly Leu Leu Leu Arg Lys Pro Asn Thr Tyr Pro Lys Met	
	360 365 370	
40	att cca gaa ttt ttc tgt gat aca tga agactttcca tatcaagaga 1265	
	Ile Pro Glu Phe Phe Cys Asp Thr	
	375 380	
45	catgggtattg actcaacagt ttccagtcac ggccaaatgt tcaatatgag tctcaataaaa 1325	
50	ctgaattttt cttgcgaaaa aaaaaaaaaa aaatccgcgg atcc 1369	
55	<210> 4	
	<211> 380	
	<212> PRT	
	<213> Homo sapiens	
	<400> 4	

EP 1 141 286 B1

	Met	Ala	Phe	Val	Cys	Leu	Ala	Ile	Gly	Cys	Leu	Tyr	Thr	Phe	Leu	Ile
	1				5					10					15	
5	Ser	Thr	Thr	Phe	Gly	Cys	Thr	Ser	Ser	Ser	Asp	Thr	Glu	Ile	Lys	Val
				20					25					30		
	Asn	Pro	Pro	Gln	Asp	Phe	Glu	Ile	Val	Asp	Pro	Gly	Tyr	Leu	Gly	Tyr
			35					40					45			
10	Leu	Tyr	Leu	Gln	Trp	Gln	Pro	Pro	Leu	Ser	Leu	Asp	His	Phe	Lys	Glu
	50						55					60				
	Cys	Thr	Val	Glu	Tyr	Glu	Leu	Lys	Tyr	Arg	Asn	Ile	Gly	Ser	Glu	Thr
15	65					70					75					80
20																
25																
30																
35																
40																
45																
50																
55																

EP 1 141 286 B1

	Trp	Lys	Thr	Ile	Ile	Thr	Lys	Asn	Leu	His	Tyr	Lys	Asp	Gly	Phe	Asp	
					85					90					95		
5	Leu	Asn	Lys	Gly	Ile	Glu	Ala	Lys	Ile	His	Thr	Leu	Leu	Pro	Trp	Gln	
				100					105					110			
	Cys	Thr	Asn	Gly	Ser	Glu	Val	Gln	Ser	Ser	Trp	Ala	Glu	Thr	Thr	Tyr	
			115					120					125				
10	Trp	Ile	Ser	Pro	Gln	Gly	Ile	Pro	Glu	Thr	Lys	Val	Gln	Asp	Met	Asp	
				130			135					140					
	Cys	Val	Tyr	Tyr	Asn	Trp	Gln	Tyr	Leu	Leu	Cys	Ser	Trp	Lys	Pro	Gly	
15	145					150					155					160	
	Ile	Gly	Val	Leu	Leu	Asp	Thr	Asn	Tyr	Asn	Leu	Phe	Tyr	Trp	Tyr	Glu	
				165					170						175		
	Gly	Leu	Asp	His	Ala	Leu	Gln	Cys	Val	Asp	Tyr	Ile	Lys	Ala	Asp	Gly	
20				180					185					190			
	Gln	Asn	Ile	Gly	Cys	Arg	Phe	Pro	Tyr	Leu	Glu	Ala	Ser	Asp	Tyr	Lys	
				195				200					205				
25	Asp	Phe	Tyr	Ile	Cys	Val	Asn	Gly	Ser	Ser	Glu	Asn	Lys	Pro	Ile	Arg	
	210						215					220					
	Ser	Ser	Tyr	Phe	Thr	Phe	Gln	Leu	Gln	Asn	Ile	Val	Lys	Pro	Leu	Pro	
	225					230					235					240	
30	Pro	Val	Tyr	Leu	Thr	Phe	Thr	Arg	Glu	Ser	Ser	Cys	Glu	Ile	Lys	Leu	
				245						250					255		
	Lys	Trp	Ser	Ile	Pro	Leu	Gly	Pro	Ile	Pro	Ala	Arg	Cys	Phe	Asp	Tyr	
				260				265						270			
35	Glu	Ile	Glu	Ile	Arg	Glu	Asp	Asp	Thr	Thr	Leu	Val	Thr	Ala	Thr	Val	
			275					280					285				
	Glu	Asn	Glu	Thr	Tyr	Thr	Leu	Lys	Thr	Thr	Asn	Glu	Thr	Arg	Gln	Leu	
40			290				295					300					
	Cys	Phe	Val	Val	Arg	Ser	Lys	Val	Asn	Ile	Tyr	Cys	Ser	Asp	Asp	Gly	
	305				310						315					320	
45	Ile	Trp	Ser	Glu	Trp	Ser	Asp	Lys	Gln	Cys	Trp	Glu	Gly	Glu	Asp	Leu	
				325						330					335		
	Ser	Lys	Lys	Thr	Leu	Leu	Arg	Phe	Trp	Leu	Pro	Phe	Gly	Phe	Ile	Leu	
				340					345					350			
50	Ile	Leu	Val	Ile	Phe	Val	Thr	Gly	Leu	Leu	Leu	Arg	Lys	Pro	Asn	Thr	
			355					360					365				
	Tyr	Pro	Lys	Met	Ile	Pro	Glu	Phe	Phe	Cys	Asp	Thr					
		370					375					380					

55
 <210> 5
 <211> 17
 <212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 5

5 ksrctccabk crtcca 17

<210> 6

<211> 20

<212> DNA

<213> Artificial Sequence

10 <220>

<223> oligonucleotide

<400> 6

atagttaaac cattgccacc 20

<210> 7

15 <211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

20 <400> 7

ctccattcgc tccaaattcc 20

<210> 8

<211> 21

<212> DNA

25 <213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 8

agtctatctt acttttactc g 21

30 <210> 9

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

35 <223> oligonucleotide

<400> 9

catctgagca ataaatattc ac 22

40 Claims

1. Use of an IL-13 antagonist in the manufacture of a medicament for treating an IL-13-related condition in a mammalian subject, wherein said antagonist is an antibody to IL-13 or an IL-13-binding fragment thereof and wherein said IL-13 related condition is atopy, allergic condition, asthma, immune complex disease, inflammatory condition of the lungs, immune deficiency, or cancer.
2. The use of Claim 1, wherein said condition is an IgE-mediated condition.
3. The use of Claims 1 or 2, wherein said condition is atopy.
4. The use of Claims 1 or 2, wherein said condition is an allergic condition.
5. The use of Claims 1 or 2, wherein said condition is asthma.
6. The use of Claims 1 or 2, wherein said condition is an immune complex disease.
7. The use of Claim 6, wherein said immune complex disease is lupus, nephritis, thyroiditis, glomerulonephritis, Grave's disease, or nephrotic syndrome.

8. The use of Claim 7, wherein said immune complex disease is lupus.
9. The use of Claim 7, wherein said immune complex disease is nephritis.
- 5 10. The use of Claim 7, wherein said immune complex disease is thyroiditis.
11. The use of Claim 7, wherein said immune complex disease is glomerulonephritis.
12. The use of Claim 7, wherein said immune complex disease is Grave's disease.
- 10 13. The use of Claim 7, wherein said immune complex disease is nephrotic syndrome.
14. The use of any one of Claims 1 to 13, wherein said antagonist is an antibody to IL-13.
- 15 15. The use of any one of Claims 1 to 13, wherein said antagonist is an IL-13-binding fragment of an IL-13 antibody.
16. Use of an IL-13 antagonist in the manufacture of a medicament for inhibiting the interaction of IL-13 with an IL-13bc protein in a condition selected from atopy, allergic condition, asthma, immune complex disease, inflammatory condition of the lungs, immune deficiency, or cancer, wherein said antagonist is an antibody to IL-13 or an IL-13-binding fragment thereof.
- 20 17. The use of Claim 16, wherein said immune complex disease is lupus, nephrotic syndrome, nephritis, thyroiditis, glomerulonephritis, or Grave's disease.
- 25 18. Use of an IL-13 antagonist in the manufacture of a medicament for treating atopy in a mammalian subject, wherein said antagonist is an antibody to IL-13 or an IL-13-binding fragment thereof.

Patentansprüche

- 30 1. Verwendung eines IL-13 Antagonisten zur Herstellung eines Arzneimittels zur Behandlung einer Erkrankung bei einem Säuger, die in Zusammenhang mit IL-13 steht, wobei dieser Antagonist ein IL-13 Antikörper oder ein an IL-13 bindendes Fragment davon ist, und wobei diese in Zusammenhang mit IL-13 stehende Erkrankung Atopie, eine allergische Erkrankung, Asthma, eine Immunkomplexerkrankung, eine entzündliche Erkrankung der Lunge, eine Immunschwäche oder Krebs ist.
- 35 2. Die Verwendung gemäß Anspruch 1, wobei diese Erkrankung eine IgE-vermittelte Erkrankung ist.
3. Die Verwendung gemäß Ansprüchen 1 oder 2, wobei diese Erkrankung Atopie ist.
- 40 4. Die Verwendung gemäß Ansprüchen 1 oder 2, wobei diese Erkrankung eine allergische Erkrankung ist.
5. Die Verwendung gemäß Ansprüchen 1 oder 2, wobei diese Erkrankung Asthma ist.
- 45 6. Die Verwendung gemäß Ansprüchen 1 oder 2, wobei diese Erkrankung eine Immunkomplexerkrankung ist.
7. Die Verwendung gemäß Anspruch 6, wobei diese Immunkomplexerkrankung Lupus, Nephritis, Thyroiditis, Glomerulonephritis, Morbus Basedow oder Nephrotisches Syndrom ist.
- 50 8. Die Verwendung gemäß Anspruch 7, wobei diese Immunkomplexerkrankung Lupus ist.
9. Die Verwendung gemäß Anspruch 7, wobei diese Immunkomplexerkrankung Nephritis ist.
10. Die Verwendung gemäß Anspruch 7, wobei diese Immunkomplexerkrankung Thyroiditis ist.
- 55 11. Die Verwendung gemäß Anspruch 7, wobei diese Immunkomplexerkrankung Glomerulonephritis ist.
12. Die Verwendung gemäß Anspruch 7, wobei diese Immunkomplexerkrankung Morbus Basedow ist.

13. Die Verwendung gemäß Anspruch 7, wobei diese Immunkomplexerkrankung Nephrotisches Syndrom ist.

14. Die Verwendung gemäß einem der Ansprüche 1 bis 13, wobei dieser Antagonist ein IL-13 Antikörper ist.

5 15. Die Verwendung gemäß einem der Ansprüche 1 bis 13, wobei dieser Antagonist ein an IL-13 bindendes Fragment eines IL-13 Antikörpers ist.

16. Verwendung eines IL-13 Antagonisten zur Herstellung eines Arzneimittels zur Hemmung der Interaktion von IL-13 mit einem IL-13bc Protein bei einer Erkrankung, die ausgewählt ist aus Atopie, einer allergischen Erkrankung, Asthma, einer Immunkomplexerkrankung, einer entzündlichen Erkrankung der Lunge, einer Immunschwäche oder Krebs, wobei dieser Antagonist ein IL-13 Antikörper oder ein an IL-13 bindendes Fragment davon ist.

17. Die Verwendung gemäß Anspruch 16, wobei diese Immunkomplexerkrankung Lupus, Nephrotisches Syndrom, Nephritis, Thyroiditis, Glomerulonephritis oder Morbus Basedow ist.

18. Die Verwendung eines IL-13 Antagonisten zur Herstellung eines Arzneimittels zur Behandlung von Atopie bei einem Säugler, wobei dieser Antagonist ein IL-13 Antikörper oder ein an IL-13 bindendes Fragment davon ist.

20 Revendications

1. Utilisation d'un antagoniste de l'IL-13 dans la fabrication d'un médicament pour traiter un état associé à l'IL-13 chez un sujet qui est un mammifère, dans laquelle ledit antagoniste est un anticorps dirigé contre l'IL-13 ou un fragment de celui-ci se fixant à l'IL-13 et dans laquelle ledit état associé à l'IL-13 est l'atopie, un état allergique, l'asthme, une maladie immunitaire complexe, un état inflammatoire des poumons, une déficience immunitaire ou le cancer.

2. Utilisation selon la revendication 1, dans laquelle ledit état est un état médié par les IgE.

3. Utilisation selon la revendication 1 ou 2, dans laquelle ledit état est l'atopie.

4. Utilisation selon la revendication 1 ou 2, dans laquelle ledit état est un état allergique.

5. Utilisation selon la revendication 1 ou 2, dans laquelle ledit état est l'asthme.

6. Utilisation selon la revendication 1 ou 2, dans laquelle ledit état est une maladie immunitaire complexe.

7. Utilisation selon la revendication 6, dans laquelle ladite maladie immunitaire complexe est le lupus, la néphrite, la thyroïdite, la glomérulonéphrite, la maladie de Graves ou le syndrome néphrotique.

8. Utilisation selon la revendication 7, dans laquelle ladite maladie immunitaire complexe est le lupus.

9. Utilisation selon la revendication 7, dans laquelle ladite maladie immunitaire complexe est la néphrite.

10. Utilisation selon la revendication 7, dans laquelle ladite maladie immunitaire complexe est la thyroïdite.

11. Utilisation selon la revendication 7, dans laquelle ladite maladie immunitaire complexe est la glomérulonéphrite.

12. Utilisation selon la revendication 7, dans laquelle ladite maladie immunitaire complexe est la maladie de Graves.

13. Utilisation selon la revendication 7, dans laquelle ladite maladie immunitaire complexe est le syndrome néphrotique.

14. Utilisation selon l'une quelconque des revendications 1 à 13, dans laquelle ledit antagoniste est un anticorps dirigé contre l'IL-13.

15. Utilisation selon l'une quelconque des revendications 1 à 13, dans laquelle ledit antagoniste est un fragment d'un anticorps dirigé contre l'IL-13 qui se fixe à l'IL-13.

16. Utilisation d'un antagoniste de l'IL-13 dans la fabrication d'un médicament pour inhiber l'interaction de l'IL-13 avec

une protéine IL-13bc dans un état choisi parmi l'atopie, un état allergique, l'asthme, une maladie immunitaire complexe, un état inflammatoire des poumons, une déficience immunitaire ou le cancer, dans laquelle ledit antagoniste est un anticorps dirigé contre l'IL-13 ou un fragment de celui-ci se fixant à l'IL-13.

5 **17.** Utilisation selon la revendication 16, dans laquelle ladite maladie immunitaire complexe est le lupus, le syndrome néphrotique, la néphrite, la thyroïdite, la glomérulonéphrite ou la maladie de Graves.

10 **18.** Utilisation d'un antagoniste de l'IL-13 dans la fabrication d'un médicament pour traiter l'atopie chez un sujet qui est un mammifère, dans laquelle ledit antagoniste est un anticorps dirigé contre l'IL-13 ou un fragment de celui-ci se fixant à l'IL-13.

15

20

25

30

35

40


45

50

55

IL-13

IL-11




IL-4

Mock

Fig. 1

FIGURE 2



sIL-13R α 2.Fc antagonizes OVA induced mucus formation

FIGURE 3

